



What is claimed is:

1. A compound represented by the following formula 1:



[1]

wherein Z is



n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-CO_2Z'$; $-CO_2R'''$; $-NH_2$; $-NHR'''$; $-NR_2'''$; $-OH$; $-OR'''$; $-CONR_2''''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;


A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-.

2. A compound represented by the following formula 1:



wherein Z is



H; A''; or B'';

n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-\text{CO}_2\text{Z}'$; $-\text{CO}_2\text{R}'''$; $-\text{NH}_2$; $-\text{NHR}'''$; $-\text{NR}_2'''$; $-\text{OH}$; $-\text{OR}'''$; $-\text{CONR}_2'''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;


A'' independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-.


3. A compound represented by the following formula 1:



wherein Z is



or



n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a , b and c represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; -CONR₂'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OR'''; -CONR₂'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;


A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-.

4. A compound represented by the following formula 1:



wherein Z is



n, m, q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;


A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;

or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and


X and X' independently represent >NH, >NR''', -O-, or -S-.

5. A compound represented by the following formula 1:



[1]

wherein Z is



H; A''; or B'';

n, m, q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;


A'' independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-.


6. A compound represented by the following formula 1:



wherein Z is



or



n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a , b and c represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-\text{CO}_2\text{Z}'$; $-\text{CO}_2\text{R}'''$; $-\text{NH}_2$; $-\text{NHR}'''$; $-\text{NR}_2'''$; $-\text{OH}$; $-\text{OR}'''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R' independently represents a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-\text{CO}_2\text{Z}'$; $-\text{CO}_2\text{R}'''$; $-\text{NH}_2$; $-\text{NHR}'''$; $-\text{NR}_2'''$; $-\text{OR}'''$; $-\text{CONR}_2'''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;

or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-.


7. The compound of claim 1 that is 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2,4-dioxothiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylic acid.

8. The compound of claim 1 that is 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylamide.


9. The compound of claim 1 that is 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5ylmethyl)-phenoxy]-phenyl}-N,N-dimethyl-acrylamide.

10. A pharmaceutical composition comprising:

a) a compound represented by the following formula 1:



wherein Z is



n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S-configuration;

R and R' each independently represent a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-\text{CO}_2Z'$; $-\text{CO}_2R'''$; $-\text{NH}_2$; $-\text{NHR}'''$; $-\text{NR}_2'''$; $-\text{OH}$; $-\text{OR}'''$; $-\text{CONR}_2'''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-; and

b) a physiologically acceptable carrier .


11. A pharmaceutical composition comprising:

a) a compound represented by the following formula 1:



[1]

wherein Z is



H; A''; or B'';

n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S-configuration;

R and R' each independently represent a hydrogen atom; linear or branched C_1 - C_{20} alkyl; linear or branched C_2 - C_{20} alkenyl; $-CO_2Z'$; $-CO_2R'''$; $-NH_2$; $-NHR'''$; $-NR_2'''$; $-OH$; $-OR'''$; $-CONR_2'''$; halogen atom; optionally substituted linear or branched C_1 - C_{20} alkyl; optionally substituted linear or branched C_2 - C_{20} alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

A'' independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-; and

b) a physiologically acceptable carrier.


12. A pharmaceutical composition comprising:

a) a compound represented by the following formula 1:




[1]

wherein Z is



or



n, m, q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a, b and c represent double bonds which may be

present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; -CONR₂''''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OR'''; -CONR₂''''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

R'''' independently represents a hydrogen atom; optionally substituted C₁-C₂₀ alkyl; optionally substituted C₁-C₂₀ alkoxy; optionally substituted C₂-C₂₀ alkenyl; optionally substituted C₆-C₁₀ aryl; or NR₂'''' represents a cyclic moiety;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-; and


b) a physiologically acceptable carrier.

13. A pharmaceutical composition comprising:

a) a compound represented by the following formula 1:



wherein Z is



n, m, q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5

provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S-configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-; and


b) a physiologically acceptable carrier.

14. A pharmaceutical composition comprising:

a) a compound represented by the following formula 1:



wherein Z is



H; A''; or B'';

n, m, q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a and b represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z configuration and, when absent, the resulting stereocenters may have the R- or S-configuration;

R and R' each independently represent a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, and A' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

A'' independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; or halo;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;


or A and B jointly, A' and B' jointly, or A'' and B'' jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR''', -O-, or -S-; and


- b) a physiologically acceptable carrier.

15. A pharmaceutical composition comprising:


- a) a compound represented by the following formula 1:



wherein Z is



or



n , m , q and r independently represent integers from zero to 4 provided that $n + m \leq 4$ and $q + r \leq 4$; p and s independently represent integers from zero to 5 provided that $p + s \leq 5$; a , b and c represent double bonds which may be present or absent; when present, the double bonds may be in the E or Z

configuration and, when absent, the resulting stereocenters may have the R- or S- configuration;

R independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OR'''; -CONR₂'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R'' independently represents a hydrogen atom; linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; -CO₂Z'; -CO₂R'''; -NH₂; -NHR'''; -NR₂'''; -OH; -OR'''; halogen atom; optionally substituted linear or branched C₁-C₂₀ alkyl; optionally substituted linear or branched C₂-C₂₀ alkenyl;

R''' independently represents a linear or branched C₁-C₂₀ alkyl; linear or branched C₂-C₂₀ alkenyl; or -(CH₂)_x-Ar, where x represents an integer from 1 to 6 and Ar represents aryl;

Z' represents a hydrogen atom or a pharmaceutically acceptable counter-ion;

A, A' and A'' each independently represent a hydrogen atom; C₁-C₂₀ acylamino; C₁-C₂₀ acyloxy; C₁-C₂₀ alkanoyl; C₁-C₂₀ alkoxycarbonyl; C₁-C₂₀ alkoxy; C₁-C₂₀ alkylamino; C₁-C₂₀ alkylcarboxylamino; carboxyl; cyano; halo; or hydroxy;

B, B' and B'' each independently represent; C₂-C₂₀ alkenoyl; aroyl; aralkanoyl; nitro; optionally substituted, linear or branched C₁-C₂₀ alkyl; or optionally substituted, linear or branched C₂-C₂₀ alkenyl;

or A and B jointly, A' and B' jointly, or A" and B" jointly, independently represent a methylenedioxy or ethylenedioxy group; and

X and X' independently represent >NH, >NR'', -O-, or -S-; and

b) a physiologically acceptable carrier.

16. The pharmaceutical composition of claim 10, wherein said compound represented by formula I is 3-(3,5-dimethoxyphenyl)-2-{4-[4-(2,4-dioxothiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylic acid.

17. The pharmaceutical composition of claim 10, wherein said compound represented by formula I is 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5-ylmethyl)-phenoxy]-phenyl}-acrylamide.

18. The pharmaceutical composition of claim 10, wherein said compound represented by formula I is 3-(3,5-dimethoxy-phenyl)-2-{4-[4-(2,4-dioxo-thiazolidin-5ylmethyl)-phenoxy]-phenyl}-N,N-dimethyl-acrylamide.